

Exam #2

Name Kex

(20) 1. Matching (use a letter only once)

The thermodynamic definition of entropy is that dS equals e.

For an increase in entropy there is an g in disorder.

The Gibbs energy G equals c.

For a spontaneous process at constant T and P , ΔG is r.

At the a point, the solid, liquid, and vapor phases are in equilibrium,

The chemical potential is the partial molar i.

For an ideal liquid or gas mixture, h or l is zero.

Raoult's Law is f.

Protein unfolding is an p process.

Henry's Law applies when the solute mole fraction approaches t.

$$c = 3.00 \times 10^8 \text{ m/s}$$

$$h = 6.626 \times 10^{-34} \text{ J s}$$

$$R = 8.314 \text{ J/mol-K}$$

$$R = 1.987 \text{ cal/mol-K}$$

$$R = 0.08206 \text{ L-atm/mol-K}$$

$$N_A = 6.02 \times 10^{23}$$

$$1 \text{ atm} = 101.325 \text{ kPa}$$

$$k_B = 1.381 \times 10^{-23} \text{ J/K}$$

a. triple

b. ΔS_{mix}

c. $H - TS$

d. one

e. dq_{rev}/T

f. $P_1 = x_1 P_1^*$

g. decrease

h. ΔH_{mix}

i. Gibbs energy

j. Tdq_{rev}

k. $P_1 = P_1^*/x_1$

l. enthalpy

m. positive

n. exothermic

o. critical

p. endothermic

q. increase

r. negative

s. $H + TS$

t. zero

(10) 2. Select the correct equation for ΔS .

Ideal gas expansion or compression: d

Heating or cooling a pure substance without a phase change at constant P: a

A phase transition at equilibrium: b

Mixing two ideal gases: e

Transition between two states with probabilities W_1 and W_2 : f

- a. $C_p \ln(T_2/T_1)$
- b. $\Delta H_{\text{trans}}/T_{\text{trans}}$
- c. $k_B \ln(W)$
- d. $nR \ln(V_2/V_1)$
- e. $-R(n_1 \ln x_1 + n_2 \ln x_2)$
- f. $k_B \ln(W_2/W_1)$
- g. $C_p(T_2 - T_1)$
- h. $k_B \ln(W_2 \times W_1)$

(10) 3. The molar heat of vaporization of ethanol is 39.3 kJ/mol and the boiling point of ethanol is 78.3 °C. Calculate ΔS_{vap} for 0.25 mole of ethanol

$$\Delta S_{\text{vap}} = \frac{39.3 \text{ kJ/mol} \times 0.25 \text{ mole}}{351.3 \text{ K}} = 0.0278 \text{ kJ/K} = 27.8 \text{ J/K}$$

unit, +1
0.25 moles, +2
°C → K, +1

eg., +6

(10) 4. Calculate the value of ΔS in heating 2 moles of an ideal gas from 50 to 100 °C at constant pressure; $C_p = 5R/2$.

$$\Delta S = C_p \ln \frac{T_2}{T_1} = \frac{5}{2} \times 8.314 \times \ln \frac{373}{323} = 5.98 \text{ J/K}$$

unit, +1
2 moles, +1
°C → K, +2

Eg., +6

(10) 5. Explain why molar S° values may be only zero or positive.

\bar{S}° at 0 K is zero for a perfect crystal and greater than zero [$\bar{S}^\circ(0\text{K}) = k_B \ln W$] for an imperfect crystal with disorder. As the temperature is increased from 0 K, the substance acquires more disorder and \bar{S}° increases!

+8 state 2nd Law

